

ANNUAL REPORT OF COOPERATIVE REGIONAL PROJECTS
Supported by Allotments of the Regional Research Fund,
Hatch Act, as Amended August 11, 1955
January 1 to December 31, 1987

- I. PROJECT: NORTH CENTRAL REGIONAL PROJECT NC-7
Introduction, Multiplication, Evaluation, Preservation,
Cataloguing, Enhancement, and Utilization of Plant Germplasm

II. COOPERATING AGENCIES AND PRINCIPALS LEADERS:

A: Administrative Adviser R. L. Mitchell, Missouri

B. Regional Coordinator R. L. Clark, Iowa

C. State Experiment Stations and Representatives

1. Alaska	*R.L. Taylor	8. Missouri	*P. Beuselinck
2. Illinois	*T. Hymowitz	9. Nebraska	*D.J. Andrews
3. Indiana	*J. Janick	10. N. Dakota	*J. Franckowiak
4. Iowa	*I. T. Carlson	11. Ohio	*S.Z. Berry
5. Kansas	*C. E. Wassom	12. S. Dakota	*A. Boe
6. Michigan	*A. Iezzoni, Chm.	13. Wisconsin	*W. Gabelman
7. Minnesota	*H. Pellett		

* Voting members

D. U. S. Department of Agriculture

1. ARS Plant Introduction Office	G. A. White
2. ARS National Program Staff, Germplasm	H. S. Shands
3. ARS Area Director, Midwest Area	G. E. Carlson
4. Cooperative State Research Service	S. Wiggans
5. Soil Conservation Service	*E. T. Jacobson
6. Northern Regional Research Center, Peoria	*R. Kleiman, Secy.
7. National Seed Storage Laboratory	S. Eberhart

E. North Central Regional Plant Introduction Station, Ames, Iowa

1. USDA-ARS staff	
a. Research Leader and Supervisory Plant Pathologist	R.L. Clark
Secretary	L. Sunstrum
b. Research Agronomist	W. W. Roath
Agricultural Research Technician	B.J. Morrell
c. Horticulturist	M. Widrlechner
d. Research Entomologist	R.L. Wilson
i. Agricultural Research Technician (Insects)	S.G. McClurg
ii. Agricultural Research Technician (Insects)	W. Hotchkiss
e. Research Plant Pathologist	Vacant

2. Iowa State University Staff

- | | |
|--|---------------|
| a. Farm Superintendent | L.L. Lockhart |
| i. Field-Lab Technician III | D. Iutjen |
| ii. Field-Lab Technician II | G. Brownrigg |
| iii. Secretary II | R. Diedrichs |
| iv. Clerk Typist II | L. Minor |
| b. Curator/Coordinator (GRIN system) and Corn Collection | M.J. Millard |
| i. Clerk IV | C. Latta |
| c. Research Associate II (Plant Pathology) | C.C. Block |
| d. Research Associate I (Grasses, Brassica) | B.C. Abel |
| e. Research Associate I (vegetables) | K. Reitsma |
| f. Research Associate I (Sunflowers, miscell. legumes) | J. Pomeroy |
| g. Predoctoral Research Assoc. (Amaranthus) | J. Lehmann |

III. PROGRESS OF WORK AND PRINCIPAL ACCOMPLISHMENTS

A. Introduction of new germplasm

New accessions entering the NC-7 Program in 1987 totaled 4,548.

B. Germplasm Multiplication

Seed increases were attempted on over 1500 accessions at Ames including: 119 Amaranthus in the field, 470 Brassicas, 201 Helianthus, 126 cucumbers, 127 corns, 107 Cuphea, 95 annual Medicago spp., 64 carrots, 53 pumpkins, 48 cantaloupes, 44 parsley, 33 Setarias, 19 parsnips, and 15 chicories. In addition, 400 annual Medicagos and 160 perennial alfalfas were increased under contract at Riverside, CA and Prosser, WA, respectively. Another 100 beets were increased in isolation field tents in Logan, UT. Leafcutter bees in cages accomplished the pollinations on the perennial Medicago spp. grown in Prosser. The annual Medics are self-pollinators so were grown without cages.

Pollination control in corn, sunflowers, and pumpkins was accomplished by hand-pollination, sibbing heterozygous lines, selfing inbreds. At least 200 plants are grown (corn and sunflowers) and attempts are made to involve all plants in the sibbing process with at least 100 seed-bearers represented in the harvest. Pumpkins are planted in 24 plant plots, 3 plants in each of 8 hills. We try to obtain at least three sibbed fruits per hill.

Wild sunflowers, cucumbers, cantaloupes, Brassicas, carrots, parsley, parsnip, chicory, and Cuphea were increased in cages. Pollinators were usually honey bees but flies were also used for the members of the carrot family. Leafcutters were the pollinators for Cuphea. We hired a full-time Technician in 1987 to manage all our insect pollinators.

The trial planting of 400 annual Medics in Riverside, CA was a success and that program will be continued in 1988. Likewise, the first year's increase of perennial Medicago spp. at Prosser, WA was very successful and will be continued. Both of these programs will be managed by the Pullman, WA, Plant Introduction Station in 1988 because we transferred maintenance responsibility for all Medicago species to that site in late 1987.

C. Germplasm Distribution

There were 295 domestic seed requests and 97 from foreign scientists. Packets distributed in response to these requests were: 12,972 and 6,216 domestic and foreign, respectively. We distributed 7,225 packets into the North Central Region, 2,651 into the Western, 2,470 into the Southern, and 563 into the North East Region. This year's total (19,188 packets) distribution is more than 3,000 higher than a year ago, indicating an increased interest in our germplasm.

By crops, this year's distribution was highest for cucumber (4,806) followed by the oilseed Brassicas (4,162), corn (2,486), sunflowers (2137), tomato (2001), alfalfa (1277), beets (642), carrots (604), and Amaranthus (372).

D. Germplasm Evaluation

1. Woody Ornamental Regional Trials.

a. Regional cooperators were sent 704 ornamental plants of 13 accessions for trial at 28 sites. An additional 128 plants of these accessions were sent to arboreta.

b. Computer-generated "Report of Planting," "Plot Information," and "First-year Performance Report" forms were distributed to trial sites this spring and the returned, completed forms are being compiled.

2. Cooperator Evaluations

a. Vegetables

Two early blight resistant tomato lines were released by North Carolina that derived their resistance from PI 126445, Lycopersicon hirsutum. The Florida variety Florida Petite had PI 205046 as one parent. The Michigan breeding population, MI-2000, released in 1987, derived its bacterial canker resistance from PI 330727 (Bulgaria 12).

Other useful traits reported in 1987 were: Fusarium resistance in Asparagus (PI 207461), Aphanomyces root rot in peas, and tolerance to the herbicide Trifluralin in Cucurbita pepo (PI 234252) and C. maxima (PI 182202).

b. Oil Seed and Special Crops

Two open-pollinated sunflower cultivars from Russia, Novinka (PI 430538) and Progress (PI 430541) were used as resistant parents in a study of the inheritance of race 3 downy mildew resistance. Various accessions of Amaranthus, Helianthus and Cuphea are being used as source materials for biotechnical studies on mechanisms responsible for production of specific chemicals.

c. Legumes

Two diploid alfalfa accessions (PI 325408 and 410970) were used as resistant parents in a study of the inheritance of Phytophthora root rot resistance. Three accessions of tetraploid alfalfa (PI 345641, 404217, and 410975), were used in the development of experimental variety NAPB32, and the latter two accessions were also involved in experimental NAPB31. Six other alfalfas are being used in a program to select a grazing-tolerant alfalfa in Georgia. Of 1200 tested for Phytophthora root rot in Wisconsin, 13 showed good resistance and will be studied further.

Two accessions of milkvetch, Astragalus cicer (PI 133142 and 362245), showed excellent vigor and winterhardiness at the Elsberry, MO Plant Materials Center. Two accessions of castor bean were used in physiologic studies as a source of glyoxysomes in a Minnesota study.

d. Grasses and Corn

Grass genera used and showing promise for various traits include: Beckmannia (forage production and quality), Eragrostis (forage production), Sorghum (greenbug resistance) and Lolium (anther culture and haploid plant regeneration). Lolium was also being used in photoperiod induction work in a grown chamber study in Missouri.

Corn accessions were reported to have: good combining ability, enhanced emergence from deep planting, drought and heat tolerance, improved protein, increased lysine, improved standability, flooding tolerance, and first generation European corn borer resistance.

3. Evaluations at the PI Station

a. Agronomic and horticultural traits

Over 500 accessions of oilseed Brassica spp. were evaluated for flowering date, seed pod arrangement, pod beak length, total pod length, pod width, locule number, pod surface, plant height, harvest date, 100-seed weight, seed color, and germination percentage of newly harvested seed. Winterhardiness data were obtained on 217 accessions of Brassica planted in six locations in North America. Data were also taken on 33 accessions of Setaria italica for heading date, leaf number and width, stem number and texture, panicle width and length, plant height and harvest date.

b. Pest evaluations

Northern corn leaf blight (race 1), common rust, and common smut evaluations were made on 400 accessions in the field. Results are being analyzed. Alternaria leaf blight evaluation techniques are under development for greenhouse screening of the sunflower collection. The cucumber collection (721 accessions) was evaluated for powdery mildew reaction in greenhouse tests. Highest resistance occurred in PI 197088, 288238, 321006, and 390258.

Silk feeding resistance to corn earworm in 107 corn accessions was evaluated in lab tests, along with 40 popcorns in field tests.

Screening for resistance to sunflower moth was carried out on 223 sunflower PI lines as part of the Sunflower CAC-approved evaluation plan.

IV. Plans for next year

We will continue to carry on an active germplasm acquisition program aimed at eliminating gaps in our collections. The most active program in 1987 was acquisition of tropical corn lines from Mexico, Colombia, and Peru. This will continue and is expected to lead to an additional 1500 lines in 1988.

We will continue to work closely with the CAC's for maize, sunflower, forage grasses, sugarbeet, cruciferous crops, leafy vegetables, root and bulb crops, and vine crops to coordinate our efforts with other parts of the National Plant Germplasm System.

We plan to acquire several hundred accessions from the collection of industrial crops housed at the USDA's Northern Regional Research Center in Peoria and enter them into our regeneration, evaluation, storage, and distribution program.

V. Publications

From the PI Station Staff

CLARK, R. L. 1987. Tomato. In: McGraw-Hill Yearbook of Science & Technology, 1988. P 451-453.

JARVIS, J. L., R. L. CLARK, C. C. BLOCK, E. C. BERRY and W. D. GUTHRIE. 1987. Interaction Between Tillage Systems, Maize Hybrids, European Corn Borers, and Stalk Rot Pathogens. *Maydica* 32:125:137.

JAWORSKI, C. A., S. C. PHATAK, S. R. GHATE, R. D. GITAITIS, and M. P. WIDRLECHNER. 1987. GA 1565-2-4 BWT, GA 219-1-2 BWT, GA 1095-1-4 BWT, and GA 1405-1-2 BWT Bacterial wilt-tolerant tomato. *HortScience* 22:324-325.

MILLER, J. F., D. C. ZIMMERMAN, B. A. VICK, and W. W. ROATH. 1987. Registration of sixteen high oleic sunflower germplasm lines and bulk population. *Crop Sci.* 27:1323.

PELLETT, H. M. and M. P. WIDRLECHNER. 1987. Landscape plant crop advisory committee. *North Central Nurserymen* 9(11):13,16-18.

PELLETT, H. M. and M. P. WIDRLECHNER. 1987. National nursery crop advisory committee - purposes and activities. *Amer. Assoc. of Bot. Gardens and Arboreta, Proc.* (In press)

ROATH, W. W. 1987. Development of new oilseed crops. *Proc. Ia. Acad. Sci.* 94:no. 4.

ROATH, W. W. 1987. Effect of day length and night temperatures on flower initiation of two *Cuphea* species. *Proc. 7th Guayule Rubber Soc. Conf.*

ROATH, W. W. and J. F. MILLER. 1987. Registration of four sunflower synthetic germplasm populations. *Crop Sci.* 27:616.

ROATH, W. W. and J. F. MILLER. 1987. Registration of HA850, HA851, HA852, and HA853 oilseed sunflower parental lines. *Crop Sci.* 27:825.

ROATH, W. W., J. F. MILLER, and T. GULYA. 1987. Registration of six oilseed sunflower parental lines. *Crop Sci.* 27:825-826.

ROATH, W. W., J. F. MILLER, and T. GULYA. 1987. Registration of three oilseed sunflower germplasm pools ND-BLPL2, ND-BLOS, and ND-RLOS. *Crop Sci.* 27:373.

ROATH, W. W. and M. P. WIDRLECHNER. 1987. Inducing germination of dormant Cuphea seed and the effects of various induction methods on seedling survival. Seed Sci. & Tech. (In press)

ROATH, W. W. and M. P. WIDRLECHNER. 1986. Sunflower pollen storage:vital stain vs. seed set. Agronomy Abstracts 1986:79.

WIDRLECHNER, M. P. 1987. Mint family as plants for honeybee forage. National Herb Growing and Marketing Conference, Proc. p. 66-76.

WIDRLECHNER, M. P. 1987. Variation in the breeding system of Lycopersicon pimpinellifolium: implications for germplasm maintenance. IBPGR Plant Genetic Resources Newsletter 70:38-43.

WILSON, R. L. 1987. Evaluation of maize plant introductions for resistance to corn earworm silk feeding. (Abstr.) N. Central Br., Entomol. Soc. Am., Submitted Papers, 1987:191.

WILSON, R. L. 1987. Evaluation of selected corn plant introductions for silk-feeding resistance to corn earworm (Lepidoptera:Noctuidae). J. Econ Entomol. 80:1048-1050.

WILSON, R. L. 1987. Iowa. Ann. Plant Resist. Insects Newsletter 13:19-21.

WILSON, R. L. and D. L. OLSON. 1987. Study of Conotrachelus seniculus life history in the laboratory and develop a method of rearing. Program Abstracts 99th Session, Ia. Acad. Sci., Symposia: Ia. Sci. Foundation No. A.

WISEMAN, B. R., and R. L. WILSON. 1987. Corn earworm development on meridic diets containing pollinated and unpollinated corn silks and on silks from two planting dates at two locations. Maydica (In press)

ATTACHMENT 1

V. B. From cooperators

ALABAMA

CASPER, R. D. and J. D. NORTON. 1985. 'Aurora' - a High Quality, Disease Resistant Cantaloupe. HortScience 20:656.

ARIZONA

LAFERRIERE, J. E. 1986. Interspecific Hybridization in Sunflowers: An Illustration of the Importance of Wild Genetic Resources in Plant Breeding. Outlook Agric. 15:104-109.

CALIFORNIA

BLAKER, N. S. and J. D. HEWITT. 1987. A Comparison of Resistance to Phytophthora parasitica in Tomato. Phytopathology 77:1113-1116.

BLAKER, N. S. and J. D. HEWITT. 1987. Comparison of Seedling and Mature Plant Resistance to Phytophthora parasitica in Tomato. HortScience 22:103-105.

CHETELAT, R. T., J. W. DEVERNA and C. M. RICK. 1987. Isozyme analysis of a Lycopersicon esculentum - Solanum lycopersicoides sesquidiploid hybrid and derived alien addition lines. TGC No. 37:39-41.

DEMPSEY, W. H. and T. H. I. SHERIF. 1987. Genetic and morphological comparisons of 7 "hairless" mutants in the tomato. TGC Reg. No. 37:43-44.

JAN, C. C. and J. M. CHANDLER. 1985. Transfer of Powdery Mildew Resistance from Helianthus debilis Nutt. to Cultivated Sunflower. Crop Science 25:664-666.

JONES, R. A. 1986. High Salt Tolerance Potential in Lycopersicon species during Germination. Euphytica 35:575-582.

KHAN, I. A., L. F. LIPPERT, and G. E. JONES. 1987. Multiple-flowering Character in Muskmelon (Cucumis melo L.). Cucurbit Genet. Coop. 10:45-46.

MCCREIGHT, J. D. 1984. Evidence of a Recessive Powdery Mildew Resistance Gene in Muskmelon PI 414723. Rep. of Cucurbit Genetic Cooperative 1984, Vol. 7:45.

MCCREIGHT, J. D., M. PITRAT, C. E. THOMAS, A. N. KISHABA and G. W. BOHN. 1987. Powdery Mildew Resistance Genes in Muskmelon. J. Amer. Soc. Hort. Sci. 112:156-160.

CALIFORNIA (Continued)

- MCCREIGHT, J. D., G. W. BOHN, and T. W. WHITAKER. 1987. PMR Honeydew Muskmelon. HortScience 22:177.
- OSBORN, T. C., D. C. ALEXANDER, and J. F. FOBES. 1987. Identification of restriction fragment length polymorphisms linked to genes controlling soluble solids content in tomato fruit. Theor. Appl. Genet. 73:350-356.
- RICK, C. M. 1987. Flowering induced in S. juglandifolium. TGC Rep. No. 37:62.
- SHANNON, M. C., J. W. GRONWALD, and M. TAL. 1987. Effects of Salinity on Growth and Accumulation of Organic and Inorganic Ions in Cultivated and Wild Tomato Species. J. Amer. Soc. Hort. Sci. 112:416-423.
- WOLF, S., D. YAKIR, M. A. STEVENS, and J. RUDICH. 1986. Cold Temperature Tolerance of Wild Tomato Species. J. Amer. Soc. Hort. Sci. 111:960-964.
- ZINK, F. W. and W. D. GUBLER. 1987. U.C. PMR45 and U.C. Top Mark Fusarium Wilt resistant (Fom-3) Muskmelon Breeding Lines. HortScience 22:172.

FLORIDA

- AUGUSTINE, J. J., B. K. HARBAUGH, and J. P. CRILL. 1981. Florida Petite: an Extremely Dwarf Tomato for Window-Sill Gardens. Circ.S. Fla. Agric. Exp. Stn., Ins. Food Agric. Sci., Univ. of Fla.

ILLINOIS

- DEAN, J. V. and J. E. HARPER. 1986. Nitric Oxide and Nitrous Oxide Production by Soybean and Winged Bean during the in Vitro Nitrate Reductase Assay. Plant Physiol. 82:718-723.
- EBERHART, S. A. 1983. Developing Virus Resistant Commercial Maize Hybrids p. 258-261. In: D. T. Gordon, J. K. Knoke, L. R. Nault and R. M. Ritter (eds) Proceedings, International Maize Virus Disease Colloquium and Workshop, 2-6 August, 1982.

KANSAS

- SKINNER, D. Z. and D. L. STUTEVILLE. 1985. Quantitatively Inherited Reactions of Alfalfa to Peronospora trifoliorum. Phytopathology 75:717-721
- SKINNER, D. Z. and D. L. STUTEVILLE. 1985. Major and Minor Genes in Diploid Alfalfa that Condition Resistance to Peronospora trifoliorum. Proc. 19th Central Alfalfa Improv. Conf. 1985, April 24-26, Oklahoma City, Ok.

KANSAS (Continued)

SKINNER, D. Z., D. L. STUTEVILLE, and D. M. STOKLI. 1986. Effects of Ancestry and Geographic Origin of Alfalfa Accessions on Resistance to Downy Mildew. Proc. 30th North Amer. Alfalfa Improv. Conf. 1986, July 27-31, St. Paul., MN

KENTUCKY

SNYDER, J. C., D. A. JOHNSON, D. E. GOOD and P. A. WESTON. 1987. Type VI trichome exudates from Chemotypes of L. hirsutum and L. hirsutum f. glabratum. TGC Rep. No. 37:67-68.

MARYLAND

BROOME, C. R., E. E. TERRELL and J. L. REVEAL. 1987. Lycopersicon esculentum P. Miller is rejected in favor of L. lycopersicum (L.) Karsten as the scientific name of the tomato. TGC Rep. No. 37:38-39.

NG, T. J. and A. N. MILLER. 1987. Genetic Analysis of Anthracnose Resistance in Tomato. (Abstr.) HortScience 21:3.

WHITE, J. L. and J. M. KAPER. 1987. Absence of Lethal Stem Necrosis in Select Lycopersicon spp. Infected by Cucumber Mosaic Virus Strain D and Its Necrogenic Satellite CARNA5. Phytopathology 77:808-811.

MICHIGAN

KAMPS, T. L., T. G. ISLEIB, R. C. HERNER, and K. C. SINK. 1987. Evaluation of Techniques to Measure Chilling Injury in Tomato. HortScience 22:1309-1312.

MISSOURI

D'ARCY, W. G. 1982. Combinations in Lycopersicon (Solanaceae). Phytologia 51:240.

NEBRASKA

POE, R. R., D. P. COYNE, B. A. SWISHER, and M. D. CLEGG. 1988. Differential Cucurbita spp. Tolerance to the Herbicide Trifluralin. J. Amer. Soc. Hort. Sci. 113-35-40

NEW MEXICO

VALLEJOS, C. E., S. D. TANKSLEY, and R. BERNATZKY. 1986. Localization in the Tomato Genome of DNA Restriction Fragments containing Sequences Homologous to the rRNA (45S), the Major Chlorophyll A/B Binding Polypeptide and the Ribulose Biphosphate Carboxylase Genes. Genetics 112:93-105.

NEW YORK

BAKER, P. B. and R. W. ROBINSON. 1985. Evaluations of Selected Cucurbita Accessions for Cucumber Beetle Complex Resistance. Search:Agriculture, 1985(31):2-8. NY Agr. Exp. Sta.

GRACEN, V. E. 1986. Sources of Temperate Maize Germplasm and Potential Usefulness in Tropical and Subtropical Environments. Advances in Agronomy 39:127-172.

PROVVIDENTI, R. 1987. Inheritance of Resistance to a Strain of Zucchini Yellow Mosaic Virus in Cucumber. HortScience 22:102-103.

NORTH CAROLINA

BOOY, G., T. C. WEHNER, and S F. JENKINS, JR. 1987. Resistance of Cucumber Lines to Rhizoctonia solani Damping-off: Not Related to Fruit Rot Resistance. HortScience 22:105-108.

GOODMAN, M. M. 1985. Exotic Maize Germplasm: Status, Prospects, and Remedies. Iowa State J. Res. 59:497-527.

NORTH DAKOTA

MILLER, J. F. and T. J. GULYA. 1987. Inheritance of Resistance to Race 3 Downy Mildew in Sunflower. Crop Sci. 27:210-212.

USDA and NORTH DAKOTA AES. 1987. Notice of Release of a Powdery Mildew Resistant Sunflower Germplasm Pool: PML. August, 1987.

OHIO

BERRY, S. Z. and G. L. OAKES. 1987. Inheritance of Resistance to Fusarium Crown and Root Rot in Tomato. HortScience 22:110-111.

OHIO (Continued)

FINDLEY, W. R., L. R. NAULT, W. E. STYER. and D. T. GORDON. 1983. Zea diploperrenis as a Source of Maize Chlorotic Dwarf Virus Resistance: A Progress Report. p. 255-257 In: D. T. Gordon, J. K. Knoke, L. R. Nault, and R. M. Ritter (eds) Proceedings International Maize Virus Disease Colloquium and Workshop, 2-6 August, 1982.

SOUTH CAROLINA

FERY, R. L. and G. G. KENNEDY. 1987. Genetic Analysis of 2-Tridecanone Concentration, Leaf Trichome Characteristics, and Tobacco Hornworm Resistance in Tomato. J. Amer. Soc. Hort. Sci. 112:886-891.

THOMAS, C. E., Y. COHEN, J. D. MCCREIGHT, E. L. JOURDAIN, and S. COHEN. 1988. Plant Disease 72:33-35.

TEXAS

DOEBLEY, J. F., M. M. GOODMAN, and C. W. STUBER. 1986. Exceptional Genetic Divergence of Northern Flint Corn. Amer. J. Bot. 73:64-69.

UTAH

HELENTJARIS, T., M. SLOCUM, S. WRIGHT, A. SCHAEFER and J. NIENHUIS. 1986. Construction of Genetic Linkage Maps in Maize and Tomato using Restriction Fragment Length Polymorphisms. Theor. Appl. Genet. 72:761-769.

VIRGINIA

O'CONNELL, M. A. and M. R. HANSON. 1985. Somatic Hybridization between Lycopersicon esculentum and Lycopersicon pennellii. Theor. Appl. Genet. 70:1-12.

WISCONSIN

HAVEY, M. J. and D. P. MAXWELL. 1987. Inheritance of Phytophthora Root Rot Resistance in Two Diploid Alfalfa Species. Crop Sci. 27:225-228.

WISCONSIN (Continued)

RENGWALSKA, M. M. and P. W. SIMON. 1986. Laboratory Evaluation of Pink Root and Fusarium Basal Rot Resistance in Garlic. Plant Disease 70:670-672.

STAUB, J. E., L. FREDRICK and T. L. MARTY. 1987. Electrophoretic variation in cross-compatible wild diploid species of Cucumis. Can. J. Bot. 65:792-798.

FOREIGN

AUSTRALIA

PATTERSON, B. D. and R. E. PAULL. 1987. Hawaii as a natural phytotron for the introgression of cold resistance. TGC Rep. No. 37:58-59.

BULGARIA

STAMOVA, L. and M. YORDANOV. 1987. Resistance to Leveillula taurica (Lev.) Arn. TGC Rep. No. 37:73.

STOEVA, P. K. and N. ZAGORSKA. 1987. Hybrids between Solanum lycopersicoides and self-compatible species from the genus Lycopersicon. TGC Rep. No. 37:73-74.

VULKOVA-ATCHKOVA, Z. N., ZAGORSKA, and E. TCHIPORIKOVA. 1987. Somaclonal changes of wild species from the genus Lycopersicon and Solanum lycopersicoides. TGC Rep. No. 37:77.

CANADA

LABERGE, C. and W. E. SACKSTON. 1987. Adaptability and Disease of Jerusalem Artichoke (Helianthus tuberosus) in Quebec. Can. J. Plant Sci. 67:349-352.

METCALF, J. G., A. A. REYES, and W. P. MOHR. 1986. Earlibright, an early Processing Tomato. Can. J. Plant Sci. 66:417-419.

POYSA, V. W. C. S. TAN, and J. A. STONE. 1987. Flooding Stress and the Root Development of Several Tomato Genotypes. HortScience 22:24-26.

COLOMBIA

VARGAS M., J. A., J. H. PAVA M., P. A. ARANGO A., and F. A. VALFLEJO C. 1984. Caracterizacion Morfo. Agronomica de Especies y Variedades Botanicas del Genero Lycopersicon. Acta Agron. 34:5-22.

ENGLAND

TAYLOR, I. B., 1986. Biosystematics of the Tomato. In: The Tomato Crop - A Scientific Basis For Improvement. J. G. Atherton and J. Rudick (Eds). Chapman and Hall, London. p. 1-34.

INDIA

BAINS, S. S. and V. PARKASH. 1985. Downy Mildew Resistance and Its Components in Muskmelon Collections. Indian Bot. Repr. 4:134-139.

BANERJEE, M. K. and KALLOO. 1987. Inheritance of Tomato Leaf Curl Virus Resistance in Lycopersicon hirsutum f. glabratum. Euphytica 36:581-584.

SREELATHAKUMARY, I., P. K. GOPALAKRISHNAN and K. V. PETER. 1987. Inheritance of combined wilt resistance in tomato. TGC Rep. No. 37:69-70.

SREELATHAKUMARY, I. and K. V. PETER. 1987. Evaluation of a set of interspecific F1 hybrids in tomato for resistance to bacterial wilt. TGC Rep. No. 37:71-72.

ISRAEL

COHEN, Y., S. COHEN, H. EYAL, and C. E. THOMAS. 1985. Inheritance of Resistance to Downy Mildew in Cucumis melo PI 124111. Cucurbit Genet. Coop. Rep. 8:36-38.

COHEN, Y., H. EYAL, J. HANANIA, and Z. MALIK. 1987. Ultrastructure of Pseudoperonospora cubensis in Muskmelons Susceptible and Resistant to Downy Mildew. (Abstr.) Phytopathology 77:1751.

KENIGSBUCH, D. and Y. COHEN. 1987. Inheritance of Resistance to Powdery Mildew Race 1 and Race 2 in Muskmelons. (Abstr.) Phytopathology 77:1724.

MILTAU, O., D. ZAMIR, and J. RUDICH. 1986. Growth Rates of Lycopersicon Species at Low Temperatures. Z. Pflanzenzuchtg. 96:193-199.

JAPAN

IMANISHI, S. 1987. Ovule culture to obtain F1 and BC1 between L. esculentum and L. peruvianum. TGC Rep. No. 37:46.

NIGERIA

KIM, S. K., Y. EFRON, F. KHADR, J. FAJEMISIN, and M. H. LEE. 1987. Registration of 16 Maize-Streak Virus Resistant Tropical Maize Parental Inbred Lines. Crop Sci. 27:824-825.

OMAN

YASSIN, T. E. 1987. Further evidence on the inheritance of resistance to leaf curl virus (LCV) disease in tomato (Lycopersicon esculentum Mill.). TGC Rep. No. 37:77-78.

YASSIN, T. E. 1985. Inheritance of resistance to leaf curl virus disease in a cross between tomato (Lycopersicon esculentum Mill.) and currant tomato (L. pimpinellifolium (Jusl.) Mill.). J. Agric. Sci., Cambridge 105:659-661.

SPAIN

AYUSO, M. C. MR. BAGUENA, J. CUARTERO and F. NUEZ. 1987. Possibilities of using the Compatible Form L. peruvianum PE-23 as a Genetic Bridge in Tomato Breeding. Tomato Genet. Coop. Rep. No. 37:36-37.

SWITZERLAND

HELLER, W. E. and C. GESSLER. 1986. Induced Systemic Resistance in Tomato Plants against Phytophthora infestans. J. Phytopathology 116:323-328.

UNKNOWN

LATERROT, H., M. BORDAT, M. RENAND, and A. MORETTI. 1987. Various levels of leafminer resistance in Lycopersicon genus. TGC Rep. No. 37:47-49.

LATERROT, H. and A. MORETTI. 1987. Extreme susceptibility of Lycopersicon cheesmanii foliage to Metribuzin herbicide. TGC Rep. No. 37-49.

UNKNOWN (Continued)

LOBO, M., A. BUSTILLO, J. I. ESCOBAR, and L. E. PETAEZ. 1987. Whitefly (Trialeurodes vaporariorum) resistance to accessions of L. pennellii. TGC Rep. No. 37:50.

LOBO, M., A. BUSTILLO, J. I. ESCOBAR, and L. E. PETAEZ. 1987. Whitefly (Trialeurodes vaporariorum) resistance in Lycopersicon hirsutum. TGC Rep. No. 37:51.

LOBO, M. and R. NAVARRO. 1987. Phoma andina resistance in interspecific hybrids between L. esculentum and L. hirsutum. TGC Rep. No. 37:52.

LOBO, M. and R. NAVARRO. 1987. Late blight horizontal resistance in L. esculentum x L. hirsutum hybrids. TGC Rep. No. 37:53.

LOBO, M., R. NAVARRO, M. M. MURILO, and A. C. RUIZ. 1987. Phoma andina resistance in Lycopersicon hirsutum accessions. TGC Rep. No. 37:53-54

APPROVED:

Amy Iezzoni, Chairman, Technical Committee

Date

R. L. Mitchell, Administrative Adviser

Date